

## Succimer is an oral metal chelator for the treatment of lead, arsenic and mercury poisoning

### Indications

- *Clinical features consistent with lead, arsenic or mercury toxicity AND a measured elevated body fluid concentration consistent with toxicity (see separate guidelines for specific indications)*
- *Succimer may have a role in at-risk asymptomatic patients with confirmed exposure to lead, arsenic or mercury (e.g. paediatric lead exposures - discuss with a clinical toxicologist)*

### Contraindications

- Known hypersensitivity
- The metal-succimer complex is eliminated via the renal tract. Consider dose adjustment with renal impairment (discuss with a clinical toxicologist)

### Presentation

Succimer 100 or 200 mg capsules (Special Access Scheme)

### Dose and administration (discuss use with a clinical toxicologist)

**Adult:** 10 mg/kg tds for 5 days, followed by 10 mg/kg bd for 14 days

### Paediatric dosing by weight:

8 – 15 kg: 100 mg tds for 5 days, followed by 100 mg bd for 14 days

16 – 23 kg: 200 mg tds for 5 days, followed by 200 mg bd for 14 days

24 - 34 kg: 300 mg tds for 5 days, followed by 300 mg bd for 14 days

35 - 44 kg: 400 mg tds for 5 days, followed by 400 mg bd for 14 days

> 45 kg: 500 mg tds for 5 days, followed by 500 mg bd for 14 days

*\*repeated courses may be needed depending on clinical features, laboratory investigations, age and co-morbidities*

### Therapeutic Endpoint:

- In general: patient asymptomatic AND laboratory investigations below concentrations indicating chelation Rx (metal concentrations should be measured 10 days after completion of chelation therapy to allow redistribution)

### Adverse effects:

- GI: Nausea, vomiting, diarrhoea, flatus, metallic taste (10-20%), mild transient transaminase rise
- Others: rash/urticaria (4%), fever, reversible neutropenia

**Pregnancy:** FDA pregnancy category C. Administration is only recommended in circumstances where potential benefit justifies the potential risk to the fetus (discuss with a clinical toxicologist).